


## How do pelvic muscles get weak?

Pelvic muscles help stop the flow of urine. Pregnancy, childbirth and being overweight can weaken the pelvic muscles. Fortunately, pelvic muscles are just like other muscles --- exercise can make them stronger. Women who leak urine may have better control of these muscles by doing pelvic muscles called Kegel exercises.

## Which muscles control my bladder?

At the bottom of the pelvis, several layers of muscles stretch between your legs. The muscles attach to the front, back and sides of the pelvic bones. Two pelvic muscles do most of the work. The biggest one stretches like a hammock. The other is shaped like a triangle (see picture below).

These are the same muscles that you would use to try to stop the flow of urine. They are the muscles you will exercise and strengthen.



## How do I exercise pelvic muscles?

You can exercise almost anywhere and any time --- while driving in a car, at your desk or watching TV. To exercise these muscles (as if you are trying to stop urine flow). Hold this squeeze for about 10 seconds, then rest for 10 seconds. Do 3 to 4 sets of 10 contractions per day.

Be patient and continue to exercise. It takes time to strengthen the pelvic muscles, just like it takes time to improve the muscles in your arms, legs or abdomen. You may not notice any changed in bladder control until after six to 12 weeks of daily exercises. Still, most women notice an improvement after just a few weeks.

## A few points to remember

- Weak pelvic muscles often lead to urine leakage.
- Daily exercises can strengthen pelvic muscles.
- These exercises often improve bladder control.
- Ask your doctor or nurse if you are squeezing the right muscles.
- Tighten your pelvic muscle before sneezing, lifting or jumping. This can prevent pelvic muscle damage and urine leakage.
- Continue to exercise. If the exercises work, continue to do them, just like any other exercises.

## Daily pelvic muscle exercise log

I exercised my pelvic muscles \_\_\_ times daily.

I spent \_\_\_ minutes exercising.

At each exercise session, I contracted my pelvic muscles \_\_\_ times.